

Effects of Intraarticular Hyaluronan on Matrix Changes Induced in the Lateral Meniscus by Total Medial Meniscectomy and Exercise

By Nigel Hope, Peter Ghosh, Thomas K.F. Taylor, Dechang Sun, and Richard Read

Total medial meniscectomy was performed in 12 adult merino sheep. Immediately after surgery, 8 animals received high-molecular-weight hyaluronan (HA) (1 mL, 10 mg/mL) and 4 were given sterile saline (1 mL) intraarticularly. Injections were given for 5 more weeks. In week 3 an exercise program, consisting of walking 24 km/wk, was initiated. This program was continued until the animals were killed at week 26 postmeniscectomy. At necropsy the lateral menisci were removed and divided into three concentric zones—inner, middle, and outer. Powdered aliquots of tissues from each zone were analyzed for collagen and hexuronate contents using colormetric methods. The glycosaminoglycans (GAGs)—chondroitin-O-sulfate (C-O-S), chondroitin-4-sulfate (C-4-S), chondroitin-6-sulfate (C-6-S), and dermatan sulfate (DS)—were determined using a high-performance liquid chromatography method. The

lateral menisci from the joints of animals injected with HA showed higher hexuronate and GAG levels than those of controls. This increase was mainly due to C-6-S, which had highest levels in the inner and middle meniscal zones. In addition, dermatan sulfate levels increased significantly in the middle and outer zones of the lateral menisci compared with the same zones of the meniscus from the saline-treated group. Collagen and C-O-S levels were not statistically different from those of controls. These data suggest that intraarticular administration of high-molecular-weight HA immediately after open total medial meniscectomy may help preserve the proteoglycans in the lateral meniscus remaining in the joint.

Copyright © 1993 by W.B. Saunders Company

INDEX WORDS: Meniscectomy; meniscus; proteoglycans; exercise; hyaluronan.

THE MENISCI (semilunar cartilages) are important structures for optimal function of the knee joint. They are shock-absorbing tissues which transmit up to 60% of the loads acting across the joint.¹⁻⁵ Apart from reducing the contact stresses on articular cartilage by increasing congruency between the curved femoral and tibial surfaces, they stabilize the joint both laterally and anterior-posteriorly.^{6,7} In flexion and extension the menisci are displaced anterior-posteriorly⁷ and their movement during extension and flexion is believed to drag synovial fluid over the articular cartilage surface, thereby facilitating its nutrition and lubrication.^{6,8}

A meniscus may be injured or torn, particularly during sporting activities; if the damaged structure interferes with normal joint function and produces pain, meniscectomy may be indicated. However, it is now recognized that meniscectomy is not a benign procedure. Total, and to a lesser extent partial, excision of a meniscus would, on the grounds discussed above, increase the contact and shear stresses acting on articular cartilage, which could lead to cartilage degeneration. Indeed, meniscectomy in rabbits,⁹⁻¹² dogs,¹³⁻¹⁶ and sheep¹⁷⁻²⁰ has been

used for many years to produce experimental models of osteoarthritis (OA). Furthermore, retrospective clinical studies²¹⁻²⁴ have demon-

From the Raymond Purves Bone and Joint Research Laboratories (University of Sydney) at the Royal North Shore Hospital of Sydney, St Leonards, New South Wales, Australia; and the School of Veterinary Studies, Murdoch University, Murdoch, Western Australia, Australia.

Supported by Pharmacia AB, Uppsala, Sweden, and the National Health and Medical Research Council of Australia.

Nigel Hope, MB, BS: Research Scholar, Raymond Purves Bone and Joint Research Laboratories (University of Sydney) at the Royal North Shore Hospital of Sydney; Peter Ghosh, PhD, FRACI, FRSC: Director, Raymond Purves Bone and Joint Research Laboratories, Royal North Shore Hospital of Sydney, Associate Professor, Department of Surgery, University of Sydney, Thomas Taylor, DPhil, FRACS, FRCS, FRCS (Ed): Professor of Orthopaedics, Royal North Shore Hospital of Sydney; Dechang Sun, MD: Research Assistant, Raymond Purves Bone and Joint Research Laboratories (University of Sydney) at the Royal North Shore Hospital of Sydney; Richard Read, PhD, FACVSc: Senior Lecturer, School of Veterinary Studies, Murdoch University.

Address reprint requests to Peter Ghosh, PhD, FRACI, FRSC, Raymond Purves Bone and Joint Research Laboratories (University of Sydney) at the Royal North Shore Hospital of Sydney, St Leonards, NSW, 2065, Australia.

*Copyright © 1993 by W.B. Saunders Company
0049-0172/93/2206-1006\$5.00/0*

strated a high incidence of OA in humans after meniscectomy.

Although the degenerative changes that occur in the matrix of articular cartilage as a consequence of meniscectomy have represented a focus of research, data on the effects of unicompartmental meniscectomy on the composition and structure of the remaining meniscus could also be important. Such knowledge could contribute to a better understanding of the remodeling response of fibrocartilage to altered mechanical loading. In addition, if the composition of the remaining meniscus is altered as a consequence of meniscectomy its ability to absorb and dissipate mechanical stresses could be modified.¹⁻⁴

Intraarticular hyaluronan (HA) has been used for more than 20 years for viscosupplementation of synovial fluid in posttraumatic equine OA and idiopathic OA in humans with reported clinical benefits.²⁵ More recently, intraarticular HA has been used in arthroscopic surgery to reduce adhesions and scar formation.²⁶⁻²⁸ Although the effects of intraarticular administration of high-molecular-weight HA on the articular cartilage of medially meniscectomized joints have been investigated,^{19,20} the influence of such treatment on the composition of the remaining lateral meniscus has not been examined. The present study was undertaken to address this question using an established meniscectomy model in sheep.¹⁷⁻²⁰

MATERIALS AND METHODS

Twelve purebred adult Merino wethers were studied. All animals underwent unilateral medial meniscectomy using a procedure described previously.¹⁷ Before meniscectomy, 8 of the 12 sheep were randomly selected to receive HA. Immediately after surgery, when the surgical opening had been sutured, each animal received intraarticularly 1.0 mL (10 mg/mL) of sterile high-molecular-weight HA (Healon; molecular weight, $\sim 3.5 \times 10^6$) preparation, which was kindly provided by Dr G. Smedegård of Pharmacia AB, Uppsala, Sweden. The remaining four meniscectomized sheep were each given 1.0 mL of sterile saline by the same route as for the HA injections.

For the next 2 weeks all animals were maintained in a communal pen (3 m \times 6 m), and each received a single weekly intraarticular injection of HA or saline. Three weeks postmeniscectomy,

when all animals had received three injections, they began an exercise regime. This consisted of walking the group (3 to 4 km/h) around a level 2-km oval track covered in loose gravel and grass. Over the following 2 weeks the number of circuits was increased until the animals walked 8 km/d, three times a week. This exercise program of 24 km/wk was maintained for the duration of the experiment, ie, 6 months postmeniscectomy. During the first 3 weeks of track work each animal received a single weekly injection of either HA or saline. Therefore, each animal received a total of six injections postmeniscectomy.

Dissection and Tissue Preparation

Animals were killed by intravenous infusion of sodium pentobarbitone (60 mg/mL) (Nembutal; Abbott Laboratories, Sydney, Australia). Lateral menisci were dissected from the meniscectomized joints, rinsed in ice-cold physiological saline, dried with a paper tissue, transferred to airtight plastic bags, and frozen until required. Before analysis each meniscus was thawed at 4°C and dissected into three longitudinal sections corresponding to the inner, middle, and outer zone of the meniscus. Each zone represents a concentric one third of the meniscus, as described previously.²⁹ Each zone was then refrozen in liquid N₂ and powdered in this medium using a Wiley Mill (A.J. Thomas, Philadelphia, PA) in which the cutting chamber had been precooled with liquid N₂. The individually powdered meniscal zones were transferred to glass containers, lyophilized, stoppered, and stored in desiccated containers at 4°C until analysis.

Analysis

Prew weighed aliquots of lyophilized powdered meniscal tissues were hydrolyzed in 6N HCl by heating at 100°C for 16 hours. The hydroxyproline released was determined using the method of Stegeman and Stalder.³⁰ The collagen content was estimated from these values by multiplying by 7.4.²⁹ Aliquots were also subjected to papain digestion³¹ and the hexuronic acid levels determined by the method of Blumenkrantz and Asboe-Hansen³² using glucuronyl lactone (Sigma Chemical Co, St Louis, MO) as a standard. The glycosaminoglycans (GAGs) present were identified and quantitated according to the high-

performance liquid chromatography (HPLC) method described below.

Sample Preparation for GAG Analysis

The method used to identify and quantitate cartilage GAGs was based on that described by Greiling et al.³³ However, modifications were made to minimize the number of sample manipulations and facilitate analysis of large numbers of samples. Aliquots (~5 mg) of lyophilized tissue were weighed into 1.5-mL Eppendorf tubes. Two hundred fifty microliters of papain digestion buffer containing 0.1 mol/L Na-acetate, 0.005 mol/L ethylenediaminetetraacetic acid (EDTA), 0.01 mol/L cysteine, and 2 μ L/mL papain suspension (Sigma Chemical Co) was added, and the tissues were hydrolyzed overnight at 60°C. One milliliter of absolute ethanol was added to the papain digest, and the GAGs precipitated overnight at 4°C on an orbital shaker at 150 rpm. The precipitate was collected by centrifugation and the supernatant decanted. The precipitated GAGs were vacuum-dried before being redissolved in 400 μ L of chondroitinase digestion buffer consisting of 0.3 mol/L Na-acetate, pH 6.5. Aliquots (180 μ L each) of enzyme digest were pipetted into 0.7-mL tapered glass crimp-top vials (Chromacol Ltd, London, England) which were compatible with a LS-3200 LC Auto-Sampler (SGE Scientific Pty Ltd, Melbourne, Australia). Ten microliters of 0.2 mol/L NaF and 10 μ L (0.05 U) each of chondroitin sulfate lyase AC or ABC (Sigma Chemical Co) were added, and the tubes were crimp sealed and incubated overnight at 37°C.

HPLC

The unsaturated disaccharides derived from the GAGs as the end product of depolymerization with either chondroitin sulfate lyase AC or ABC were fractionated by HPLC and spectrophotometrically detected at 232 nm using a semiautomated HPLC system. This consisted of a Dynamax NH2-column (4.6 \times 250 mm) fitted with a guard column (4.6 \times 15 mm) (Rainin Instrument Co Inc, Woburn, MA) that was eluted with an aqueous buffer consisting of 0.3 mol/L ammonium acetate and 20 mmol/L glycine, pH 5.5, at a flow rate of 1 mL/min provided by a ETP/Kortec K25M single-piston pump (ICI Instru-

ments, Rydalmere, NSW, Australia). The ultraviolet detector was an ERMA model ERC-7210 (ERMA Optical Works Ltd, Tokyo, Japan), and signal analysis and peak integration were performed using a SMAD/SMAD chromV.2.13 system (Morgan Kennedy Research, Rose Bay, NSW, Australia) controlled by an Apple Macintosh computer.

Pure standards of HA (Healon; Pharmacia [Australia] Pty Ltd, North Ryde, NSW, Australia), chondroitin-4-sulfate (C-4-S) (Calbiochem, Alexandria, NSW, Australia), and disc chondroitin-6-sulfate (C-6-S) donated by Professor Pearce, University of British Columbia, Vancouver, Canada, and dermatan sulfate (DS) (Sigma Chemical Co) were used for the quantitation of GAGs. The amount of DS present in each GAG precipitate was calculated by the difference of 4-sulfated disaccharide values obtained by chondroitin sulfate lyase ABC and AC digestion.

Methods of Statistical Analysis

Because four control and eight HA-injected animals were used for this study, analytical data were analyzed using the unpaired two-tailed *t* test. Differences between HA- and saline-treated groups were considered statistically significant at $P \leq .05$.

RESULTS

At necropsy a partially regrown meniscus replica was found in the medial compartment of all animals. This structure was integrated with the synovium from which it probably originated.¹⁵ No significant differences were observed between the experimental groups in terms of the size or mass of these regrown tissues (data not shown). As noted previously,²⁹ the outer region of the menisci contained more collagen than the inner, but differences were not observed between the levels of this protein in the same zones of menisci from placebo (saline)- or HA-injected joints (data not shown). In contrast, the hexuronate values of the inner and middle zones of the lateral menisci derived from HA-injected joints were 50% to 100% higher than those from the same zones of menisci from the placebo-injected animals ($P < .05$) (Fig 1). This elevation in hexuronate values was also mirrored in total GAG levels as deter-




Fig 1: Hexuronate content ($\mu\text{g}/\text{mg}$ dry weight of tissue) of the various zones of lateral menisci from meniscectomized joints of animals injected intra-articularly with saline (placebo) or HA (mean \pm SE). Meniscal zones were defined as inner (\square), middle ($\▤$), outer ($\▥$), and pooled (\blacksquare). Statistically significant differences ($*P < .05$) existed between the groups for the inner, middle, and pooled zones.

mined by HPLC, but for these analyses statistical significance ($P < .05$) was demonstrable between the groups only for data pooled from all meniscus regions (Fig 2).

The comparison of individual GAGs of the menisci of the two groups showed no difference in their chondroitin-0-sulfate (present mainly as HA) content (Fig 3), but marked differences in the other chondroitin sulfate isomers were observed. C-4-S levels were elevated in the outer meniscal zone of the HA-injected group compared with the same zone of the placebo-treated group ($P < .05$) (Fig 4). C-6-S levels were higher in the same tissues as C-4-S, but again in the middle region of the menisci from the HA-injected animals (Fig 5). DS, although less abundant than the other chondroitin sulfate isomers, was elevated in the middle and outer zones of the lateral menisci of the HA-injected joints compared with the same zones of the menisci in the placebo group ($P < .05$) (Fig 6).

DISCUSSION

The knee joint menisci are fibrocartilaginous tissues whose composition and structure are well

adapted for their functions. Because they are wedge-shaped in cross-section and the coefficient of friction of synovial fluid is low, they are squeezed outwardly when the joint is loaded. Radial displacement of the menisci is resisted by their tibial attachments, and a circumferential "hoop strain" develops in the periphery of the structure on joint loading.^{4,34} To accommodate these high tensile stresses, the outer rim of the meniscus is composed predominantly of thick collagen fibers aligned circumferentially. The inner and middle zones of the meniscus, while still rich in collagen, contain much higher amounts of proteoglycans than the outer zone.^{15,29,35} The composite structure of the meniscus is thus well suited to withstand and recover from both compressional and tensional stresses.^{15,36} Biochemically and morphologically,^{29,35,37} the inner and middle zones of the meniscus show many similarities to hyaline articular cartilage except that the proteoglycan content is lower and the collagen present is largely type I rather than type II.³⁵

Studies by Tribe³⁸ and Squire et al³⁹ on grazing patterns of sheep have shown that on average these animals traverse 4 km/d in fenced paddocks




Fig 2: Total GAG content ($\mu\text{g}/\text{mg}$ dry weight of tissue) of various zones of lateral menisci from meniscectomized joints of animals injected intra-articularly with saline (placebo) or HA (mean \pm SE). Meniscal zones were defined as inner (\square), middle ($\▤$), outer ($\▥$), and pooled (\blacksquare). Statistically significant differences ($*P < .05$) existed between the groups for the pooled zones.




Fig 3: Chondroitin-O-sulfate (mainly HA) content ($\mu\text{g}/\text{mg}$ dry weight of tissue) of lateral menisci from meniscectomized joints of animals injected intra-articularly with saline (placebo) or HA (mean \pm SE). Meniscal zones were defined as inner (\square), middle ($\▤$), outer ($\▨$), and pooled (\blacksquare).

irrespective of the area. The exercise regimen used in the present experiments, in which sheep were walked 24 km/wk around a grass track, is therefore comparable with their normal activity pattern, bearing in mind that some movements also occurred while the animals were penned.

In a previous study⁴⁰ it was shown that the hexuronate contents of all zones of both the medial and lateral menisci of unoperated sheep subjected to an exercise program similar to that used in the present experiments were elevated compared with the same zones of a nonexercised, nonoperated group. This finding shows that the fibrochondrocytes of the knee joint menisci, like the chondrocytes of articular cartilage,^{41,42} can adapt to a new mechanical environment by modifying biosynthetic activities to elaborate an extracellular matrix more suited to the types of stresses to which they are subjected.

The most notable difference between the outcomes for the two postoperative regimens used in the present experiments was the higher concentration of proteoglycans in all zones of the lateral menisci of the HA-injected group (Figs 1 and 2). This increase could be interpreted, on the

basis of previous studies,⁴⁰⁻⁴² to indicate that in the HA-treated group the lateral menisci were subjected to higher dynamic compressional stresses than the corresponding menisci of the saline-treated group. The finding that C-6-S was the major contributor to the elevated hexuronate values, particularly in the middle zone (Fig 5), was compatible with this reasoning.^{41,42} Increased loading of the lateral menisci after 6 weekly injections of high-molecular-weight HA could arise from improved weight bearing on the meniscectomized joint because force-plate studies using the same ovine model used here, but with HA preparations of lower molecular weight, showed that lameness was diminished in the actively treated groups compared with saline controls.^{18,19}

On the other hand, the immediate postoperative injection of HA into the meniscectomized joints may have preserved meniscal proteoglycan levels directly. High-molecular-weight HA is reported to encourage healing of experimentally created defects in rabbit anterior cruciate ligaments.⁴³ However, the rate of healing of circular defects in rabbit menisci after a single injection of HA into the joints was found to be indistin-




Fig 4: C-4-S content ($\mu\text{g}/\text{mg}$ dry weight of tissue) of the various zones of lateral menisci from meniscectomized joints of animals injected intra-articularly with saline (placebo) or HA (mean \pm SE). Meniscal zones were defined as inner (\square), middle ($\▤$), outer ($\▨$), and pooled (\blacksquare). Statistically significant differences ($*P < .05$) existed between the outer zones of the two groups.




Fig 5: C-6-S content ($\mu\text{g}/\text{mg}$ dry weight of tissue) of the various zones of lateral menisci from meniscectomized joints of animals injected intraarticularly with saline (placebo) or HA (mean \pm SE). Meniscal zones were defined as inner (\square), middle ($\▤$), outer ($\▨$), and pooled (\blacksquare). Statistically significant differences ($*P < .05$) existed between groups for the middle and pooled zones.

guishable from that of saline controls.⁴⁴ The reduced proteoglycan contents of the lateral menisci removed from saline treated joints could have initially arisen as a consequence of enhanced matrix catabolism. This may have resulted from the postoperative trauma and local inflammation associated with open meniscectomy. In this regard the documented antiinflammatory properties of HA,⁴⁵⁻⁴⁸ coupled with its ability to attenuate oxygen-derived free radical⁴⁹ and proteolytic⁴⁹⁻⁵² breakdown of cartilage may have provided some protection to meniscal proteoglycans.

The C-6-S and C-4-S of human^{35,37} and canine³⁵ menisci have been analyzed and would appear to be present in these tissues as large aggregating proteoglycans (aggrecans) similar to those isolated from hyaline cartilage.^{35,37} Indeed, they may be encoded from the same gene.³⁷ Similarly, the elevated levels of DS in the middle and outer zones of the menisci of the HA-treated group presumably arise from an increase in DS-containing proteoglycans (DS-PGs). Two DS-PGs, decorin and biglycan, have been isolated from human menisci.³⁷ Biglycan consists of a core

protein of molecular weight $\sim 42,500$ to which two DS chains (molecular weight $\sim 37,000$) are attached. Decorin has a different core protein to biglycan and has only one DS chain. The DS chains for both DS-PGs are hybrids containing iduronic and glucuronic acid residues glycosidically linked mainly to 4-sulfated *N*-acetyl galactosamine (see reviews^{53,54} for collected references).

Although the biological role of the DS-PGs is still the subject of research, it is clear that they have many functions. It has been shown that decorin is associated with the d band of type I and II collagen fibrils and can inhibit fibrillogenesis in vitro.⁵⁴ Cartilage-derived decorin and biglycan have been shown to inhibit attachment and spreading of 3T3 cells to a fibronectin-coated matrix.^{53,55} However, because neither the DS chains nor the core proteins alone could inhibit cell attachment, it was concluded that the DS-PG complex was necessary for inhibition. These studies have led to the proposal that the DS-PGs bind to a specific receptor on fibronectin adjacent to the integrin-binding site. When DS-PGs are




Fig 6: DS content ($\mu\text{g}/\text{mg}$ dry weight of tissue) of the various zones of lateral menisci from meniscectomized joints of animals injected intraarticularly with saline (placebo) or HA (mean \pm SE). Meniscal zones were defined as inner (\square), middle ($\▤$), outer ($\▨$), and pooled (\blacksquare). Statistically significant differences ($*P < .05$) existed between groups for the middle and outer zones.

bound to this receptor, the DS side chains sterically block the integrin-binding receptor, thereby inhibiting cell adhesion. Additional interactions between the GAG chains and fibronectin may augment this process.⁵³ Decorin can also bind transforming growth factor β (TGF- β) via its core-protein⁵⁶ and modulate the activities of this ubiquitous growth factor. This has led to the suggestion that decorin may be a regulator of cell proliferation. However, TGF- β also stimulated the synthesis of biglycan and decorin by fibroblast and ovary cells,^{56,57} and because DS-PGs can bind TGF- β a negative-feedback regulation of cell mitosis has been proposed.⁵⁶ These and other⁵⁸ experiments have led to the suggestion that the DS-PGs inhibit connective tissue repair processes.⁵³

Dynamic loading (5 kg/0.3 Hz) of bovine sesamoid cartilage in vitro stimulates the biosynthesis of DS-PGs, especially decorin.⁵⁹ Similar results have been obtained for degenerative articular cartilage from adult sheep joints subjected

to lateral meniscectomy.⁶⁰ The deposition of DS-PGs in hyaline cartilage therefore would seem to be associated with chondrocyte response to high dynamic loading and presumably represents part of an adaptive reorganization of the extracellular matrix in response to mechanical overload. Whether the enrichment of the lateral menisci of the HA-treated joints with DS-PGs is also related to altered mechanical loading of this tissue or arises from a direct pharmacological effect of HA has yet to be determined. However, the short half-life of HA in synovial joints^{61,62} suggests that either or both of these mechanisms would provide the maximum benefit during the immediate postoperative period, when tissue remodeling and repair are most active.

ACKNOWLEDGMENT

The authors thankfully acknowledge the advice and assistance of D. Burkhardt in the HPLC experiments and thank L. Campbell for typing the manuscript.

REFERENCES

1. Seedhom BB, Hargreaves DJ: Transmission of the load in the knee joint with special reference to the role of the menisci. *Eng Med* 8:220-228, 1979
2. Kurosawa H, Fukubayashi T, Nakajima H: Load bearing mode of the knee joint: Physical behaviour of the knee joint with or without menisci. *Clin Orthop* 149:283-290, 1980
3. Krause WR, Pope MH, Johnson RJ, et al: Mechanical changes in the knee after meniscectomy. *J Bone Joint Surg* 59:599-604, 1986
4. Ahmed AM: The load-bearing role of the menisci, in Mow VC, Arnoczky SP, Jackson DW (eds): *Knee Meniscus Basic and Clinical Foundations*. New York, NY, Raven, 1992, pp 59-73
5. Walker PS, Erkman MJ: The role of the menisci in force transmissions across the knee. *Clin Orthop Rel Res* 109:184-192, 1975
6. Smillie IS: *Injuries of the Knee Joint* (ed 3). Baltimore, MD, Williams & Wilkins, 1962
7. Levy IM, Torzilli PA, Fisch ID: The contribution of the menisci to the stability of the knee, in Mow VC, Arnoczky SP, Jackson DW (eds): *Knee Meniscus Basic and Clinical Foundations*. New York, NY, Raven, 1992, pp 107-115
8. MacConaill MA: The function of intra-articular fibrocartilages with special reference to the knee and inferior radio-ulna joints. *J Anat* 66:210-227, 1931
9. Floman Y, Eyre DR, Glimcher MJ: Induction of osteoarthritis in the rabbit knee joints: Biochemical studies on the articular cartilage. *Clin Orthop Rel Res* 147:278-286, 1980
10. Shapiro F, Glimcher MJ: Induction of osteoarthritis in the rabbit knee joint: Histologic changes following meniscectomy and meniscal lesions. *Clin Orthop Rel Res* 147:287-295, 1980
11. Moskowitz R, Goldberg V, Malesud C: Metabolic responses of cartilage in experimentally induced osteoarthritis. *Ann Rheum Dis* 40:584-592, 1981
12. Colombo C, Butler M, O'Byrne E, et al: Development of knee joint pathology following lateral meniscectomy and section of the fibular collateral and sesamoid ligaments. *Arthritis Rheum* 26:875-886, 1983
13. Cox JS, Cordell LD: The degenerative effects of medial meniscus tears in dogs' knees. *Clin Orthop Rel Res* 125:236-242, 1977
14. Ghosh P, Sutherland JM, Taylor TKF, et al: The effect of postoperative joint immobilisation of articular cartilage degeneration following meniscectomy. *J Surg Res* 35:461-473, 1983
15. Ghosh P, Taylor TKF, Pettit GD, et al: Effect of postoperative immobilisation on the regrowth of the knee joint semilunar cartilage: An experimental study. *J Orthop Res* 1:153-164, 1983
16. Hannan N, Ghosh P, Bellenger C, et al: The systemic administration of glycosaminoglycan polysulphate (Arteparon) provided partial protection of articular cartilage from damage produced by meniscectomy in the canine. *J Orthop Res* 5:47-59, 1987
17. Ghosh P, Sutherland J, Bellenger C, et al: The influence of weight-bearing exercise on articular cartilage of meniscectomised joints. *Clin Orthop Rel Res* 252:101-113, 1990
18. Ghosh P, Numata Y, Smith S, et al: Animal models of early osteoarthritis and their use for the evaluation of potential chondroprotective agents. *Agents Actions Suppl* 39:195-206, 1993
19. Ghosh P, Read R, Armstrong S, et al: The effects of intra-articular administration of hyaluronan in a model of

early osteoarthritis in sheep. I. Gait analysis and radiological and morphological studies. *Semin Arthritis Rheum* 22:18-30, 1993 (suppl 1)

20. Ghosh P, Read R, Numata Y, et al: The effects of intra-articular administration of hyaluronan in a model of early osteoarthritis in sheep. II. Cartilage composition and proteoglycan metabolism. *Semin Arthritis Rheum* 22:31-42, 1993 (suppl 1)

21. Fairbank TK: Knee joint changes after meniscectomy. *J Bone Joint Surg* 39:664-670, 1948

22. Tapper EM, Hoover NW: Late results after meniscectomy. *J Bone Joint Surg [Am]* 51:517-526, 1969

23. Appel H: Late results after meniscectomy in the knee joint. A clinical and roentgenological follow-up. *Acta Orthop Scand* 133:89-99, 1970

24. Johnson RJ, Kettelkamp DB, Clark W, et al: Factors affecting late results after meniscectomy. *J Bone Joint Surg [Am]* 56:719-729, 1974

25. Ghosh P, Smith M, Wells C: Second line agents in osteoarthritis, in Furst D, Dixon J (eds): *Second Line Agents in the Treatment of Rheumatic Diseases*. New York, NY, Marcel Dekker, 1991, pp 363-427

26. Balazs EA, Denlinger JL: The role of hyaluronic acid in arthritis and its therapeutic use, in Peyron JG (ed): *Osteoarthritis: Current Clinical and Fundamental Problems*. Proceedings of a Workshop Held in Paris, 9-11 April 1984. Basle, Switzerland, Geigy, 1984, pp 165-174

27. Balazs EA, Denlinger JL: Clinical use of hyaluronan. *Ciba Found Symp* 143:267-285, 1989

28. McCain JP, Balazs EA, de la Rua H: Preliminary studies on the use of a viscoelastic solution in arthroscopic surgery of the temporomandibular joint. *J Oral Maxillofac Surg* 47:1161-1168, 1989

29. Ingman AM, Ghosh P, Taylor TKF: Variation of collagenous and non-collagenous proteins of human knee joint menisci with age and degeneration. *Gerontologia* 20:212-223, 1974

30. Stegeman H, Stalder K: Determination of hydroxyproline. *Clin Chim Acta* 189:267-273, 1967

31. Pearce RH, Mathieson JM: Quantitative isolation of purified acidic glycosaminoglycans from rat skin. *Can J Biochem* 45:1565-1576, 1976

32. Blumenkrantz N, Asboe-Hansen G: New method for quantitative determination of uronic acids. *Anal Biochem* 54:484-489, 1973

33. Greiling H, Stuhlsatz HW, Tillmanns U: Chondroitin, chondroitin 4-sulphate, chondroitin 6-sulphate, and dermatan sulphate. *Methods Enzym Anal* 6:60-69, 1984

34. Aspden RM: A model for the function and failure of the meniscus. *Eng Med* 14:119-122, 1985

35. Adams ME, Hukins DWL: The extracellular matrix of the meniscus, in Mow VC, Arnoczky SP, Jackson DW (eds): *Knee Joint, Basic and Clinical Foundation*. New York, NY, Raven, 1992, pp 15-36

36. Aspden RM, Yarker YE, Hukins DWL: Collagen orientations in the meniscus of the knee joint. *J Anat* 140:371-380, 1985

37. Roughley PJ, White RJ: The dermatan sulphate proteoglycan of the adult human meniscus. *J Orthop Res* 10:631-637, 1992

38. Tribe DE: Some seasonal observations on the behaviour of sheep. *J Agric Sci Camb* 39:309-315, 1949

39. Squire VR, Wilson AD, Daws GT: Comparison of the walking activity of some Australian sheep. *Proc Aust Soc Animal Prod* 9:376, 1972

40. Hope N, Ghosh P, Burkhardt D, et al: The effects of moderate exercise on the collagen and glycosaminoglycans of the knee joint meniscus (semi-lunar cartilage). (submitted)

41. Kiviranta I, Jurvelin J, Tammi M, et al: Weight-bearing controls glycosaminoglycan concentration and articular cartilage thickness in the knee joint of young beagle dogs. *Arthritis Rheum* 30:801-809, 1987

42. Kiviranta I, Tammi M, Jurvelin J, et al: Moderate running exercise augments glycosaminoglycans and thickness of articular cartilage in the knee joint of young beagle dogs. *J Orthop Res* 6:188-195, 1988

43. Wiig ME, Amiel D, VandeBerg J, et al: The early effect of high molecular weight hyaluronan (hyaluronic acid) on anterior cruciate ligament healing: An experimental study in rabbits. *J Orthop Res* 8:425-434, 1990

44. Hope NGA, Collier S, Ghosh P, et al: Repair of tears of the knee joint meniscus: Evaluation of the effects of hyaluronic acid using a rabbit model. Proceedings of the Australian Orthopaedic Association Annual Meeting, September 1990

45. Häkansson L, Hällgren R, Venge P: Regulation of granulocyte function by hyaluronic acid. In vitro effects on phagocytosis, locomotion and metabolism. *J Clin Invest* 66:298-305, 1984

46. Miyazaki K, Goto S, Okawara H, et al: Sodium hyaluronate (SPH): Studies on analgesic and anti-inflammatory effects of sodium hyaluronate (SPH). *Pharmacometrics* 28:1123-1135, 1984

47. Punzi L, Schiavon F, Cavasin F, et al: The influence of intra-articular hyaluronic acid on PGE₂ and cAMP of synovial fluid. *Clin Exp Rheumatol* 7:247-250, 1989

48. Tobetto K, Yasui T, Ando T, et al: Inhibitory effects of hyaluronan on [¹⁴C]arachidonic acid release from labeled human synovial fibroblasts. *Jpn J Pharmacol* 60:79-84, 1992

49. Larson NE, Lombard KM, Parent EG, et al: Effects of Hylan on cartilage and chondrocyte cultures. *J Orthop Res* 10:23-32, 1992

50. Pisko EJ, Turner RA, Soderham LP, et al: Inhibition of neutrophil phagocytosis and enzyme release by hyaluronic acid. *Clin Exp Rheumatol* 1:41-52, 1984

51. Iwata H, Miyauchi S, Machida A, et al: The effect of sodium hyaluronate on the articular cartilage treated with papain. Transactions of the Combined Meeting of the Orthopaedic Research Societies of USA, Japan and Canada, 1991, p 143

52. Kitoh Y, Katsuramaki T, Tanaka H, et al: Effect of SL-1010 (sodium hyaluronate with high molecular weight) on experimental osteoarthritis induced by intra-articularly applied papain in rabbits. *Folia Pharmacol Jpn* 100:67-76, 1992

53. Rosenberg L: Structure and function of dermatan sulphate proteoglycans in articular cartilage, in Kuettner R, Schleyerbach R, Peyron J, et al (eds): *Articular Cartilage and Osteoarthritis*. New York, NY, Raven, 1991, pp 45-62

54. Stancescu V: The small proteoglycans of cartilage matrix. *Semin Arthritis Rheum* 20:51-64, 1990

55. Lewandowska K, Choi H, Rosenberg L, et al: Fibronectin-mediated adhesion of fibroblasts—Inhibition by dermatan sulfate proteoglycan and evidence for a cryptic glycosaminoglycan-binding domain. *J Cell Biol* 105:1443-1454, 1987
56. Yamaguchi Y, Mann D, Ruoslahti E: Negative regulation of transforming growth factor-beta by the proteoglycan decorin. *Nature* 346:281-284, 1990
57. Westergren-Thorsson G, Antonsson P, et al: The synthesis of a family of structurally related proteoglycans in fibroblasts is differently regulated by TGF-beta. *Matrix* 11:177-183, 1991
58. Tollefsen DM, Peacock ME, Monafo WJ: Molecular size of dermatan sulphate oligosaccharides required to bind and activate heparin cofactor II. *J Biol Chem* 261:8854-8858, 1986
59. Visser NA, Van Kampen GPJ, Vande Stadt RJ, et al: Modulation of the synthesis of dermatan sulphate proteoglycans I and II in intact articular cartilage in vitro. *Trans Orthop Res Soc* 16:364, 1991
60. Little CB, Ghosh P, Bellenger CR: Meniscectomy increases aggrecan, decorin (PG-S2) and biglycan (PG-S1) metabolism in cartilage. *Trans Orthop Res Soc* 18:707, 1993 (abstr)
61. Brown TJ, Laurent UBG, Fraser JRE: Turnover of hyaluronan in synovial joints: Elimination of labelled hyaluronan from the knee joint of the rabbit. *Exp Physiol* 76:125-134, 1991
62. Fraser JRE, Kimpton WG, Pierseonek BK, et al: The kinetics of hyaluronan in normal and acutely inflamed synovial joints. *SemiArthritis Rheum* 22:9-17, 1993 (suppl 1)